

COLLAPSING OF CALABI-YAU MANIFOLDS AND SPECIAL LAGRANGIAN SUBMANIFOLDS

YUGUANG ZHANG

ABSTRACT. In this paper, the relationship between the existence of special lagrangian submanifolds and the collapsing of Calabi-Yau manifolds is studied. First, special lagrangian fibrations are constructed on some regions of bounded curvature and sufficiently collapsed in Ricci-flat Calabi-Yau manifolds. Then, in the opposite direction, it is shown that the existence of special lagrangian submanifolds with small volume implies the collapsing of some regions in the ambient Calabi-Yau manifolds.

1. INTRODUCTION

The notion of special lagrangian submanifold was introduced by Harvey and Lawson in the seminar paper [27]. Mclean studied the deformation theory of special lagrangian submanifolds in [34]. In the pioneer work [43], Stominger, Yau and Zaslow propose a conjecture about constructing the mirror manifold of a given Calabi-Yau manifold, the SYZ conjecture, via special lagrangian fibrations. Since then, lots of works were devoted to study special lagrangian submanifolds and fibrations (c.f. [28], [37], [38], [39], [19], [20], [33], [21], [42], [45], [30], [31], and references in [31]). In [32] and [24], a refined version of SYZ conjecture was proposed by using the collapsing of Ricci-flat Calabi-Yau manifolds in the Gromov-Hausdorff sense. These two versions of SYZ conjecture suggest a relationship between the existence of special lagrangian submanifolds and the collapsing of Calabi-Yau manifolds. In this paper, we study this relationship.

If (M, ω, J, g) is a compact Ricci-flat Kähler n -manifold, and admits a no-where vanishing holomorphic n -form Ω , the holomorphic volume form, $(M, \omega, J, g, \Omega)$ is called a Ricci-flat Calabi-Yau n -manifold, and (ω, J, g, Ω) is called a Calabi-Yau structure on M . We can normalize Ω such that

$$\frac{\omega^n}{n!} = \frac{(-1)^{\frac{n^2}{2}}}{2^n} \Omega \wedge \bar{\Omega},$$

(c.f. [31]). Yau's theorem of Calabi conjecture guarantees the existence of Ricci-flat Kähler metrics on Kähler manifolds with trivial canonical bundle (c.f. [46]), which implies the existence of Calabi-Yau structures on such manifolds. The holonomy group of a Ricci-flat Calabi-Yau n -manifold is a subgroup of $SU(n)$. The study of Calabi-Yau manifolds is important in both mathematics and physics (c.f. [47]).

A special lagrangian submanifold L of phase $\theta \in \mathbb{R}$ in a Ricci-flat Calabi-Yau n -manifold $(M, \omega, J, g, \Omega)$ is a lagrangian submanifold $L \subset M$ corresponding to the Kähler form ω such that $\operatorname{Re} e^{\sqrt{-1}\theta} \Omega|_L = dv_{g|_L}$ where $dv_{g|_L}$ denotes the volume form of $g|_L$ on L . Equivalently, $\dim_{\mathbb{R}} L = n$,

$$\omega|_L \equiv 0, \quad \operatorname{Im} e^{\sqrt{-1}\theta} \Omega|_L \equiv 0$$

Partially supported by the National Natural Science Foundation of China 09221010054.

(c.f. [27]). In [34], Mclean showed that, for a compact special lagrangian submanifold L in a Calabi-Yau manifold $(M, \omega, J, g, \Omega)$, the local moduli space of special lagrangian submanifolds near L is a smooth manifold of dimension $b_1(L)$, and, moreover, the tangent space of the moduli space at L can be identified with the space of harmonic 1-forms on $(L, g|_L)$. In [28], various structures on the moduli space of special lagrangian submanifolds were studied.

A special lagrangian fibration on a Calabi-Yau n -manifold (M, ω, Ω) consists of a topological space B , and a surjection $f : M \rightarrow B$ such that there is an open dense subset $B_0 \subset B$, which is a real n -manifold, satisfying that, for any $b \in B_0$, $f^{-1}(b)$ is a smooth special lagrangian submanifold in (M, ω, Ω) . By [14] (see also [22]), $f^{-1}(b)$, $b \in B_0$, is a n -torus. The first step of SYZ conjecture is to construct such fibration on a Calabi-Yau manifold when the complex structure is close to the large complex structure limit point enough (c.f. [43]). Then the mirror manifold is a compactification of the dual fibration of $f : f^{-1}(B_0) \rightarrow B_0$. Generalized special lagrangian fibrations were constructed in some almost Calabi-Yau manifolds in [37], [38], [39], [20]. In [23], special lagrangian fibrations were constructed on some Borcea-Voisin type Calabi-Yau 3-manifolds with degenerated Ricci-flat Kähler-Einstein metrics. In [40], H-minimal Lagrangian fibrations, a generalization of special lagrangian fibration, were constructed on some regions of Kähler-Einstein manifolds with negative scalar curvature.

In [32] and [24], SYZ conjecture was refined to the following form: Let $\pi : \mathcal{M} \rightarrow \Delta$ be a maximally unipotent degeneration of Calabi-Yau n -manifolds over the unit disc $\Delta \subset \mathbb{C}$, and α be an ample class on $\mathcal{M}$. For any $t \in \Delta \setminus \{0\}$, let $\tilde{g}_t$ be the unique Ricci-flat Kähler metric on $M_t = \pi^{-1}(t)$ with its Kähler form $\tilde{\omega}_t \in \alpha|_{M_t} \in H^{1,1}(M_t, \mathbb{R}) \cap H^2(M_t, \mathbb{Z})$, and $\tilde{g}_t = \text{diam}_{\tilde{g}_t}^{-2}(M) \tilde{g}_t$. Then $(M_t, \tilde{g}_t)$ converges to a compact metric space (B, d_B) of Hausdorff dimension n in the Gromov-Hausdorff sense, when $t \rightarrow 0$. Furthermore, there is a closed subset $S_B \subset B$ of Hausdorff dimension $n - 2$ such that $B \setminus S_B$ is an affine manifold, and d_B is induced by a Monge-Ampère metric g_B on $B \setminus S_B$ (c.f. [32]). The mirror manifolds are supposed to be constructed from the dual affine structure on $B \setminus S_B$ and the metric g_B . This conjecture was verified for some K3 surfaces in [24]. By using hyperKähler rotation, some K3 surfaces admit special lagrangian fibrations. It was shown that some K3 surfaces with Ricci-flat Kähler metrics collapse along such special lagrangian fibrations in [24]. The two versions of SYZ conjecture suggest the equivalence between the existence of special lagrangian submanifolds and the collapsing of Ricci-flat Kähler metrics on some regions of Calabi-Yau manifolds, when complex structures are close to the large complex limit point enough.

In Riemannian geometry, the collapsing of Riemannian manifolds was studied by various authors (c.f. [10], [11], [9], [13], [15], and references in [15]), since Gromov introduced the notion of Gromov-Hausdorff topology in [18]. In [11], it was proved that there is a constant $\epsilon_0(n) > 0$ depending only on n such that there is an F -structure of positive rank on the region M_{ϵ_0} in a Riemannian n -manifold (M, g) , where M_{ϵ_0} denotes the subset with injectivity radius $i_g(p) < \epsilon_0$ and sectional curvature $\sup_{B_g(p,1)} |K_g| \leq 1$, for any $p \in M_{\epsilon_0}$. See

[10] and [11] for the definition of F -structure of positive rank, which is a generalization of fibration. A folklore conjecture says that there should be special lagrangian fibrations on such region in a Calabi-Yau manifold, i.e. the region of bounded curvature and sufficiently collapsed (c.f. [16]). The first result in the present paper is devoted to construct special lagrangian fibrations under such Riemannian geometric conditions.

Theorem 1.1. *For any $n \in \mathbb{N}$ and any $\sigma > 1$, there exists a constant $\epsilon = \epsilon(n, \sigma) > 0$ depending only on n and σ such that, if $(M, \omega, J, g, \Omega)$ is a closed Ricci-flat Calabi-Yau n -manifold with $[\omega] \in H^2(M, \mathbb{Z})$, and $p \in M$ such that*

i) *the injectivity radius and the sectional curvature*

$$i_g(p) < \epsilon, \quad \sup_{B_g(p, 1)} |K_g| \leq 1,$$

ii) $[\Omega|_{B_g(p, \sigma i_g(p))}] \neq 0$ in $H^n(B_g(p, \sigma i_g(p)), \mathbb{C})$,

then there is an open subset $W \subset M$ satisfying that $B_g(p, \sigma i_g(p)) \subset W$, and (W, ω, Ω) admits a special lagrangian fibration of a phase $\theta \in \mathbb{R}$, i.e. there is a topological space B , and a surjection $f : W \longrightarrow B$ such that, for any $b \in B$, $f^{-1}(b)$ is a smooth n -submanifold,

$$\omega|_{f^{-1}(b)} \equiv 0, \quad \text{and} \quad \text{Im} e^{\sqrt{-1}\theta} \Omega|_{f^{-1}(b)} \equiv 0.$$

Remark 1.2. From the proof of this theorem, we can see that B is an orbifold, and, if b belongs the singular set of B , $f^{-1}(b)$ is a smooth multi-fiber.

Remark 1.3. The condition ii) in the theorem can be replaced by the following small non-vanishing n -cycle condition: there is an $[A] \in H_n(B_g(p, \sigma i_g(p)), \mathbb{Z})$ such that

$$\int_A \Omega \neq 0.$$

This condition can not be removed since it is satisfied if there is a special lagrangian submanifold L near p having comparable size to $i_g(p)$, for example $L \subset B_g(p, \sigma i_g(p))$.

Remark 1.4. It is a challenging task to verify condition i) in Theorem 1.1, i.e. to find the region of bounded curvature in a Ricci-flat Calabi-Yau manifold. If $(M, \omega, J, g, \Omega)$ is a K3-surface with Ricci-flat metric, it was shown in [13] that there are universal constants $C > 0$, $\tau > 0$, and a finite subset $\{p_j\} \subset M$, $1 \leq j \leq \tau$, such that

$$\sup_{B_g(p, 1)} |K_g| \leq C,$$

for any $p \in M \setminus \bigcup_{1 \leq j \leq \tau} B_g(p_j, 2)$. From the author's knowledge, no such estimate for higher dimensional Calabi-Yau manifolds is known except some trivial cases, for example $K3 \times T^2$.

Next, in the opposite direction, we show that the existence of special lagrangian submanifolds with small volume implies the collapsing of some regions in the ambient Calabi-Yau manifolds. The following theorem is a corollary of a volume comparison theorem for calibrated submanifolds in [20].

Theorem 1.5. *Let $(M, \omega, J, g, \Omega)$ be a closed Ricci-flat Calabi-Yau n -manifold, and $p \in M$. Assume that the sectional curvature K_g satisfies*

$$\sup_{B_g(p, 2\pi)} K_g \leq 1,$$

and there is a special lagrangian submanifold L of phase θ such that $p \in L$, and

$$\int_L \text{Re} e^{\sqrt{-1}\theta} \Omega < \frac{\pi}{2n} \varpi_{n-1},$$

where ϖ_{n-1} denotes the volume of S^{n-1} with the standard metric of constant curvature 1. Then the injectivity radius $i_g(p)$ of (M, g) at p satisfies that

$$i_g(p)^n \leq \frac{n\pi^{n-1}}{2^{n-1}\varpi_{n-1}} \int_L \operatorname{Re} e^{\sqrt{-1}\theta} \Omega.$$

As mentioned in Remark 1.4, the assumption of bounded curvature is not desirable. If we strengthen the condition of the existence of a special lagrangian submanifold to the existence of a special lagrangian fibration, we can still obtain the collapsing result in the absence of bounded curvature.

Theorem 1.6. *For any $n \in \mathbb{N}$ and any $\varepsilon > 0$, there is a constant $\delta = \delta(n, \varepsilon) > 0$ satisfying that: Assume that $(M, \omega, J, g, \Omega)$ is a closed Ricci-flat Calabi-Yau n -manifold, $p \in M$, and there is a homology class $A \in H_n(M, \mathbb{Z})$ such that, for any $x \in B_g(p, 1)$, there is a special lagrangian submanifold L_x of phase θ passing x and presenting A , i.e. $x \in L_x$ and $[L_x] = A$. If*

$$\int_A \operatorname{Re} e^{\sqrt{-1}\theta} \Omega < \delta,$$

then

$$\operatorname{Vol}_g(B_g(p, 1)) \leq \varepsilon.$$

Let $\{(M_k, \omega_k, J_k, g_k, \Omega_k)\}$ be a family of closed Ricci-flat Calabi-Yau n -manifolds, and $\{p_k\}$ be a sequence of points such that there are open subsets $W_k \supset B_{g_k}(p_k, 1)$ in M_k admitting special lagrangian fibrations $f_k : W_k \rightarrow B_k$ of phase θ_k corresponding to Calabi-Yau structures (ω_k, Ω_k) . If

$$\lim_{k \rightarrow \infty} \int_{f_k^{-1}(b_k)} \operatorname{Re} e^{\sqrt{-1}\theta_k} \Omega_k = 0,$$

where $b_k \in B_k$, Theorem 1.6 implies that

$$\lim_{k \rightarrow \infty} \operatorname{Vol}_{g_k}(B_{g_k}(p_k, 1)) = 0,$$

and, by passing to a subsequence, $\{(M_k, g_k, p_k)\}$ converges to a path metric space of lower Hausdorff dimension in the pointed Gromov-Hausdorff sense (c.f. [13], [4]). Theorem 1.1, Theorem 1.5 and Theorem 1.6 give an evidence of the equivalence between the existence of special lagrangian submanifolds and the collapsing of Ricci-flat Kähler metrics on Calabi-Yau manifolds near the large complex limit point from the Riemannian geometry's point of view.

Theorem 1.5 and Theorem 1.6 are special cases of corresponding theorems for Riemannian manifolds with calibration forms and calibrated submanifolds (See Section 6 for details). Special lagrangian submanifolds in Calabi-Yau manifolds are examples of calibrated submanifolds. There are some other examples of calibrated submanifolds, for example associative and coassociative submanifolds in G_2 -manifolds and Cayley submanifolds in $Spin(7)$ -manifolds, which have deep relationships with M-theory and exceptional mirror symmetry (c.f. [1] [26]). It is interesting to understand the interaction between the collapsing of G_2 -manifolds (resp. $Spin(7)$ -manifolds) and the existence of associative and coassociative submanifolds (resp. Cayley submanifolds).

The organization of the paper is as follows: In §2, we review some notions and results, which will be used in this paper. In §3, we use the blow-up argument to give local approximations of Calabi-Yau manifolds by complete flat Calabi-Yau manifolds. In §4, we study the deformation of special lagrangian fibrations. In §5, we prove Theorem 1.1 by combining the results in §3 and §4. Finally, we prove Theorem 1.5 and Theorem 1.6 in §6.

Acknowledgement: The author would like to thank Prof. Weidong Ruan and Prof. Xiaochun Rong for useful discussions. Thanks also goes to Prof. Fuquan Fang for constantly support.

2. PRELIMINARIES

In this section, we review some notions and results, which will be used in the proof of Theorem 1.1, Theorem 1.5 and Theorem 1.6.

2.1. Cheeger-Gromov convergence. Since Gromov introduced the concept of Gromov-Hausdorff convergence in [18], the convergence of Riemannian manifolds was studied from various perspectives (c.f. [2], [3], [12], [13], [15], [17], [24], [41], [44] and references in [5]). There is an extension of Gromov-Hausdorff convergence to sequences of pointed metric spaces for dealing with non-compact situations.

Definition 2.1 ([18], [15]). For two pointed complete metric spaces (X, d_X, x) and (Y, d_Y, y) , a map $\psi : (X, x) \rightarrow (Y, y)$ is called an ϵ -pointed approximation if $\psi(B_{d_X}(x, \epsilon^{-1})) \subset B_{d_Y}(y, \epsilon^{-1})$, $\psi(x) = y$, and $\psi|_{B_{d_X}(x, \epsilon^{-1})} : B_{d_X}(x, \epsilon^{-1}) \rightarrow B_{d_Y}(y, \epsilon^{-1})$ is an ϵ -approximation, i.e. $B_{d_Y}(y, \epsilon^{-1}) \subset \{y' \in Y | d_Y(y', \psi(B_{d_X}(x, \epsilon^{-1}))) < \epsilon\}$, and

$$|d_X(x_1, x_2) - d_Y(\psi(x_1), \psi(x_2))| < \epsilon$$

for any x_1 and $x_2 \in B_{d_X}(x, \epsilon^{-1})$. The number

$$d_{GH}((X, d_X, x), (Y, d_Y, y)) = \inf \left\{ \epsilon \left| \begin{array}{l} \text{There are } \epsilon - \text{pointed approximations} \\ \psi : (X, x) \rightarrow (Y, y), \text{ and } \phi : (Y, y) \rightarrow (X, x) \end{array} \right. \right\}$$

is called pointed Gromov-Hausdorff distance between (X, d_X, x) and (Y, d_Y, y) . We say that a family of pointed complete metric spaces (X_k, d_{X_k}, x_k) converges to a complete metric space (Y, d_Y, y) in the pointed Gromov-Hausdorff sense, if

$$\lim_{k \rightarrow \infty} d_{GH}((X_k, d_{X_k}, x_k), (Y, d_Y, y)) = 0.$$

The following is the famous Gromov pre-compactness theorem:

Theorem 2.2 ([18]). *Let $\{(M_k, g_k, p_k)\}$ be a family of pointed complete Riemannian manifolds such that Ricci curvatures $\text{Ric}(g_k) \geq -C$ for a constant C independent of k . Then, a subsequence of (M_k, g_k, p_k) converges to a pointed complete path metric space (Y, d_Y, y) in the pointed Gromov-Hausdorff sense.*

This theorem shows that a family of pointed complete Ricci-flat Einstein manifolds converges to a pointed complete path metric space by passing to a subsequence in the pointed Gromov-Hausdorff sense. The structure of the limit space was studied in [6], [13], [8] and [12] etc. We need the following result in the proof of Theorem 1.6.

Theorem 2.3 ([6], [5]). *Let $\{(M_k, g_k, p_k)\}$ be a family of pointed complete Ricci-flat Einstein n -manifolds, i.e. $\text{Ric}(g_k) \equiv 0$, such that*

$$\text{Vol}_{g_k}(B_{g_k}(p_k, 1)) \geq C,$$

for a constant C in-dependent of k , and (Y, d_Y, y) be a pointed complete path metric space such that

$$\lim_{k \rightarrow \infty} d_{GH}((M_k, g_k, p_k), (Y, d_Y, y)) = 0.$$

Then the Hausdorff dimension $\dim_{\mathcal{H}} Y = n$, and there is a closed subset $S_Y \subset Y$ of Hausdorff dimension $\dim_{\mathcal{H}} S_Y < n - 1$ such that $Y \setminus S_Y$ is a n -manifold, and d_Y is induced by a Ricci-flat Einstein metric g_{∞} on $Y \setminus S_Y$. Furthermore, for any compact subset $D \subset Y \setminus S_Y$, there are embeddings $F_{k,D} : D \rightarrow M_k$ such that $F_{k,D}^* g_k$ converges to g_{∞} in the C^{∞} -sense.

In [18] and [17], a convergence theorem, the Cheeger-Gromov convergence theorem, was proved for Riemannian manifolds with bounded curvature and non-collapsing. The Kähler version of this theorem can be found in [36]. See [12] for the convergence of manifolds with other holonomy groups.

Theorem 2.4 (Kähler version of Cheeger-Gromov convergence theorem). *Let $\{(M_k, g_k, J_k, \omega_k, p_k)\}$ be a family of pointed compact Kähler n -manifolds with sectional curvature and injectivity radius at p_k*

$$|K_{g_k}| \leq 1, \quad i_{g_k}(p_k) \geq C,$$

for a constant $C > 0$ independent of k . Then a subsequence of $\{(M_k, g_k, J_k, \omega_k, p_k)\}$ converges to a complete Kähler n -manifold (X, g, J, ω, p) in the pointed $C^{1,\alpha}$ -sense, i.e. for any $r > 0$, there are embeddings $F_{k,r} : B_g(p, r) \rightarrow M_k$ such that $F_{k,r}(p) = p_k$, $F_{k,r}^ g_k$ (resp. $dF_{k,r}^{-1} J_k dF_{k,r}$ and $F_{k,r}^* \omega_k$) converges to g (resp. J and ω) in the $C^{1,\alpha}$ -sense.*

If we assume that g_k are Einstein metrics, it is shown in [2] that, by passing to a subsequence, $\{(M_k, g_k, J_k, \omega_k, p_k)\}$ converges to (X, g, J, ω, p) in the pointed C^{∞} -sense, and g is also an Einstein metric, i.e. $F_{k,r}^* g_k$ (resp. $dF_{k,r}^{-1} J_k dF_{k,r}$ and $F_{k,r}^* \omega_k$) converges to g (resp. J and ω) in the C^{∞} -sense. Assume that $(M_k, g_k, J_k, \omega_k, p_k)$ are Ricci-flat Calabi-Yau manifolds, and Ω_k are the corresponding holomorphic volume forms. Since Ω_k are parallel, i.e. $\nabla^{g_k} \Omega_k \equiv 0$, for any $r > 0$, $F_{k,r}^* \Omega_k$ converge to a holomorphic volume form Ω on X in the C^{∞} -sense, and $(X, g, J, \omega, \Omega)$ is a complete Ricci-flat Calabi-Yau n -manifold.

In [10], [11], the collapsing of Riemannian manifolds with bounded curvature was studied by combining blow-up arguments and the Cheeger-Gromov convergence theorem. It was shown that there is a constant $\epsilon_0(n) > 0$ depending only on n such that there is an F -structure $\mathcal{F}$ of positive rank on a region covering M_{ϵ_0} in a Riemannian n -manifold (M, g) , where M_{ϵ_0} denotes the subset with injectivity radius $i_g(p) < \epsilon_0$ and sectional curvature $\sup_{B_g(p,1)} |K_g| \leq 1$, for any $p \in M_{\epsilon_0}$. See [10] and [11] for the definition of F -

structure of positive rank. If we assume that g is a Kähler metric, some additional information about the F -structure $\mathcal{F}$ is expected. We have the following conjecture:

Conjecture 2.5. *For any $n \in \mathbb{N}$, there exists a constant $\epsilon = \epsilon(n) > 0$ depending only on n such that, if (M, ω, J, g) is a closed Kähler n -manifold with $[\omega] \in H^2(M, \mathbb{Z})$, and*

$$M_{\epsilon} = \{p \in M \mid i_g(p) < \epsilon, \sup_{B_g(p,1)} |K_g| \leq 1\},$$

then there is an open subset $W \subset M$ such that $W \supset M_{\epsilon}$, and W admits an F -structure $\mathcal{F}$ of positive rank, whose orbits $\mathcal{O}_p$, $p \in M_{\epsilon}$, are isotropic submanifolds of (M, ω) , i.e.

$$\omega|_{\mathcal{O}_p} \equiv 0.$$

We will address this question in other papers. In the present paper, we prove Theorem 1.1 by combining Theorem 2.4 and the deformation theory of special lagrangian fibrations.

2.2. A comparison theorem for calibrated submanifolds. In [27], Harvey and Lawson introduced the notion of calibrated submanifold. If (M, g) is a Riemannian manifold, and Θ is a closed n -form such that $\Theta|_\xi \leq dv_\xi$ for any oriented n -plane ξ in the tangent bundle of M , then Θ is called a calibration on M , where dv_ξ denotes the volume form on ξ . An oriented n -submanifold L of M is called calibrated by the calibration Θ , if $\Theta|_L$ equals to the volume form of $g|_L$ on L . Mclean studied the deformation theory of calibrated submanifolds in [34].

There are some examples of calibrated submanifolds: holomorphic submanifolds in Kähler manifolds, special lagrangian submanifolds in Calabi-Yau manifolds, associative coassociative submanifolds in G_2 -manifolds, and Cayley submanifolds in $Spin(7)$ -manifolds (c.f. [27] [29]) etc. If (M, ω, J, g) is a Kähler m -manifold, then $\frac{1}{n!}\omega^n$, $n \leq m$, are calibrations on M , and holomorphic n -submanifolds are calibrated by $\frac{1}{n!}\omega^n$. If $(M, \omega, J, g, \Omega)$ is a Ricci-flat Calabi-Yau n -manifold, then, for any $\theta \in \mathbb{R}$, $\text{Re}e^{\sqrt{-1}\theta}\Omega$ is a calibration on M , and a special lagrangian submanifold L of phase θ is calibrated by $\text{Re}e^{\sqrt{-1}\theta}\Omega$. If (M, g) is a Riemannian manifold with holonomy group G_2 , then M admits a parallel 3-form ϕ , which is a calibration on M , and $*_g\phi$ is a calibration 4-form on M . Submanifolds calibrated by ϕ are called associative submanifolds, and submanifolds calibrated by $*_g\phi$ are called coassociative submanifolds (c.f. [29]). If (M, g) is a Riemannian manifold with holonomy group $Spin(7)$, then M admits a calibration 4-form Ω , and Cayley submanifolds are submanifolds calibrated by Ω (c.f. [29]).

In [20], a volume comparison theorem for calibrated submanifolds was obtained.

Theorem 2.6 (Theorem 2.0.1. in [20]). *Let (M, g) be a closed Riemannian manifold, Θ be a calibration n -form, and $p \in M$. Assume that the sectional curvature K_g satisfies*

$$\sup_{B_g(p, \frac{2\pi}{\sqrt{\Lambda}})} K_g \leq \Lambda, \quad \Lambda > 0,$$

and there is a submanifold L calibrated by Θ such that $p \in L$. Then

$$\text{Vol}_g(B_g(p, r) \cap L) \geq \text{Vol}_{h_1}(B_{h_1}(r)),$$

for any $r \leq \min\{i_g(p), \frac{\pi}{\sqrt{\Lambda}}\}$, where h_1 denotes the standard metric on S^n with constant curvature Λ , and $B_{h_1}(r)$ denotes a metric r -ball in S^n .

2.3. Implicit function theorem. For studying the deformation of special lagrangian fibrations, we need the following quantity version of implicit function theorem.

Theorem 2.7 (Theorem 3.2 in [37]). *Let $(\mathfrak{B}_1, \|\cdot\|_1)$ and $(\mathfrak{B}_2, \|\cdot\|_2)$ be two Banach spaces, $\|\cdot\|_E$ be the standard Euclidean metric on $\mathbb{R}^n$, $U \subset \mathbb{R}^n \times \mathfrak{B}_1$ be an open set, and $\mathfrak{F} : U \rightarrow \mathfrak{B}_2$ be a continuously differentiable map. Denote the differential*

$$D\mathfrak{F}(y, \sigma)(\dot{y} + \dot{\sigma}) = D_y\mathfrak{F}(y, \sigma)\dot{y} + D_\sigma\mathfrak{F}(y, \sigma)\dot{\sigma},$$

for $(y, \sigma) \in U$, $\dot{y} \in \mathbb{R}^n$ and $\dot{\sigma} \in \mathfrak{B}_1$. Assume that $(0, 0) \in U$ satisfies that $D_\sigma\mathfrak{F}(0, 0) : \mathfrak{B}_1 \rightarrow \mathfrak{B}_2$ has a bounded linear inverse $D_\sigma\mathfrak{F}(0, 0)^{-1} : \mathfrak{B}_2 \rightarrow \mathfrak{B}_1$ with

$$\|D_\sigma\mathfrak{F}(0, 0)^{-1}\| \leq \overline{C}$$

for a constant $\overline{C} > 0$. Let $r > 0$, $\delta_0 > \delta > 0$ be constants such that, if $\|y_0\|_E < r$, and $\|\sigma\|_1 \leq \delta_0$, then $(y_0, \sigma) \in U$, and

$$\|D_\sigma\mathfrak{F}(y_0, \sigma) - D_\sigma\mathfrak{F}(0, 0)\| \leq \frac{1}{2\overline{C}} \quad \text{and} \quad \|\mathfrak{F}(y_0, 0)\|_2 \leq \frac{\delta}{4\overline{C}}.$$

Then, for any $\|y\|_E < r$, there exists a unique $\sigma(y) \in \mathfrak{B}_1$ such that

$$\mathfrak{F}(y, \sigma(y)) = 0, \quad \|\sigma(y)\|_1 \leq \delta.$$

Furthermore,

$$D\sigma(y)\dot{y} = -D_\sigma \mathfrak{F}(y, \sigma)^{-1} D_y \mathfrak{F}(y, \sigma)\dot{y}.$$

The difference between this version of implicit function theorem and the usual one (c.f. [25]) is that we use the condition $\|\mathfrak{F}(y, 0)\|_2 \leq \frac{\delta}{4C}$ to replace the condition $\mathfrak{F}(0, 0) = 0$ besides other quantity estimates.

3. THE BLOW-UP LIMIT

Let $\{(M_k, \omega_k, J_k, g_k, \Omega_k)\}$ be a family of closed Ricci-flat Calabi-Yau n -manifold with $[\omega_k] \in H^2(M_k, \mathbb{Z})$, and $p_k \in M_k$. Assume that

i) the injectivity radius and the sectional curvature

$$i_{g_k}(p_k) < \frac{1}{k}, \quad \sup_{B_{g_k}(p_k, 1)} |K_{g_k}| \leq 1,$$

ii) there is a $\sigma \gg 1$ such that $[\Omega_k|_{B_{g_k}(p_k, \sigma i_{g_k}(p_k))}] \neq 0$ in $H^n(B_{g_k}(x_k, \sigma i_{g_k}(p_k)), \mathbb{C})$.

If we denote $\tilde{\omega}_k = i_{g_k}^{-2}(p_k)\omega_k$, $\tilde{g}_k = i_{g_k}^{-2}(p_k)g_k$, and $\tilde{\Omega}_k = i_{g_k}^{-n}(p_k)\Omega_k$, then

$$i_{\tilde{g}_k}(p_k) = 1, \quad \sup_{B_{\tilde{g}_k}(p_k, k)} |K_{\tilde{g}_k}| \leq \frac{1}{k^2},$$

and $[\tilde{\Omega}_k|_{B_{\tilde{g}_k}(p_k, \sigma)}] \neq 0$ in $H^n(B_{\tilde{g}_k}(p_k, \sigma), \mathbb{C})$. By the Cheeger-Gromov's convergence theorem (c.f. Theorem 2.4), a subsequence of $(M_k, \tilde{\omega}_k, \tilde{g}_k, J_k, \tilde{\Omega}_k, p_k)$ converges to a complete flat Calabi-Yau n -manifold $(X, \omega_0, g_0, J_0, \Omega_0, p_0)$ in the C^∞ -sense, i.e. for any $r > \sigma$, there are embeddings $F_{r,k} : B_{g_0}(p_0, r) \rightarrow M_k$ such that $F_{r,k}(p_0) = p_k$, and $F_{r,k}^* \tilde{g}_k$ (resp. $F_{r,k}^* \tilde{\omega}_k$ and $F_{r,k}^* \tilde{\Omega}_k$) converges to g_0 (resp. ω_0 and Ω_0) in the C^∞ -sense. The purpose of this section is to prove that (X, ω_0, Ω_0) admits a special lagrangian fibration.

By the smooth convergence, $i_{g_0}(p_0) = \lim_{k \rightarrow \infty} i_{\tilde{g}_k}(p_k) = 1$. The soul theorem (c.f. [11], [35]) implies that there is a compact flat totally geodesic submanifold $S \subset X$, the soul, such that (X, g_0) is isometric to the total space of the normal bundle $\nu(S)$ with a metric induced by $g_0|_S$ and a natural flat connection.

Lemma 3.1. $\dim_{\mathbb{R}} S \geq n$.

Proof. If $\dim_{\mathbb{R}} S < n$, then

$$H^n(X, \mathbb{C}) = H^n(T_r(S), \mathbb{C}) = H^n(S, \mathbb{C}) = \{0\}$$

for any $r > 0$, where $T_r(S) = \{p \in X | \text{dist}_{g_0}(p, S) \leq r\}$. Let $r_0 > r_1 > \sigma$ such that $T_{r_1}(S) \subset B_{g_0}(x_0, r_0)$, and $F_{r_0,k}(T_{r_1}(S)) \supset B_{\tilde{g}_k}(x_k, \sigma)$ for $k \gg 1$. Then the inclusion maps induce homeomorphisms on cohomology groups

$$H^n(M, \mathbb{C}) \rightarrow H^n(F_{r_0,k}(T_{r_1}(S)), \mathbb{C}) \rightarrow H^n(B_{\tilde{g}_k}(x_k, \sigma), \mathbb{C}),$$

$$\text{and } [\tilde{\Omega}_k] \mapsto [\tilde{\Omega}_k|_{F_{r_0,k}(T_{r_1}(S))}] \mapsto [\tilde{\Omega}_k|_{B_{\tilde{g}_k}(x_k, \sigma)}] \neq 0.$$

Thus $[\tilde{\Omega}_k|_{F_{r_0,k}(T_{r_1}(S))}] \neq 0$ in $H^n(F_{r_0,k}(T_{r_1}(S)), \mathbb{C})$, which contradicts to

$$H^n(F_{r_0,k}(T_{r_1}(S)), \mathbb{C}) \cong H^n(T_{r_1}(S), \mathbb{C}) = \{0\}.$$

□

If $\pi_h : \tilde{S} \rightarrow S$ is the holonomy covering of S , Bieberbach's theorem (c.f. [11], [35]) says that $(\tilde{S}, \pi_h^* g_0)$ is isometric to a flat torus, and π_h has finite order at most $\lambda(n)$, for a constant $\lambda(n)$ depending only on n . If we denote $\bar{\pi} : \mathbb{C}^n \rightarrow X$ the universal covering of X with $\bar{\pi}(0) \in S$, then $\bar{S} = \bar{\pi}^{-1}(S)$ is a real linear subspace of $\mathbb{C}^n$, and $\omega_E = \bar{\pi}^* \omega_0$ (resp. $\Omega_E = \bar{\pi}^* \Omega_0$) is the standard flat Kähler form (resp. the standard holomorphic volume form), i.e. $\omega_E = \sqrt{-1} \sum_{\alpha} dz_{\alpha} \wedge d\bar{z}_{\alpha}$ and $\Omega_E = dz_1 \wedge \cdots \wedge dz_n$ under some coordinates $z_1, \dots, z_n$ on $\mathbb{C}^n$. Note that there is a lattice $\Lambda \subset \bar{S}$ such that $\tilde{S} = \bar{S}/\Lambda$. If we denote $\mathfrak{q} : \bar{S} \rightarrow \tilde{S}$ the quotient map, then $\bar{\pi} = \pi_h \circ \mathfrak{q}$.

Lemma 3.2. $\dim_{\mathbb{R}} \bar{S} = n$, and there is a constant $\theta_0 \in \mathbb{R}$ such that $\omega_E|_{\bar{S}} = 0$ and $\text{Im} e^{\sqrt{-1}\theta_0} \Omega_E|_{\bar{S}} = 0$. Moreover, S is a special lagrangian submanifold of phase θ_0 in (X, ω_0, Ω_0) , i.e. $\dim_{\mathbb{R}} S = n$,

$$\omega_0|_S \equiv 0, \quad \text{and} \quad \text{Im} e^{\sqrt{-1}\theta_0} \Omega_0|_S = 0.$$

Proof. If $\omega_E|_{\bar{S}} \neq 0$, and thus $\omega_0|_S \neq 0$, then there are two vectors $v_1, v_2 \in \bar{S}$ such that $\omega_E(v_1, v_2) > 0$. By perturbing v_1 and v_2 a little bit if necessary, we have that $\tilde{\Sigma} = \mathfrak{q}(\{t_1 v_1 + t_2 v_2 | t_i \in \mathbb{R}\})$ is a closed 2-torus in $\tilde{S}$, i.e. a closed 2-parameters subgroup. Thus $\Sigma = \pi_h(\tilde{\Sigma})$ is a closed oriented surface in S , which satisfies

$$\int_{\Sigma} \omega_0 \geq \frac{1}{\lambda(n)} \int_{\tilde{\Sigma}} \pi_h^* \omega_0 \geq \frac{\omega_E(v_1, v_2)}{\lambda(n) \|v_1 \wedge v_2\|_{h_E}} V_E > 0,$$

where V_E denotes the Euclidean area of the intersection of $\{t_1 v_1 + t_2 v_2 | t_i \in \mathbb{R}\}$ with the fundamental domain of the quotient map $\mathfrak{q}$. From the smooth convergence of $(M_k, \tilde{\omega}_k, \tilde{g}_k)$,

$$\lim_{k \rightarrow \infty} i_{g_k}^{-2}(p_k) \int_{F_{r,k}(\Sigma)} \omega_k = \lim_{k \rightarrow \infty} \int_{F_{r,k}(\Sigma)} \tilde{\omega}_k = \lim_{k \rightarrow \infty} \int_{\Sigma} F_{r,k}^* \tilde{\omega}_k = \int_{\Sigma} \omega_0,$$

for $r \gg 1$ such that $\Sigma \subset B_{g_0}(p_0, r)$. Thus

$$0 < \frac{1}{2} i_{g_k}^{-2}(p_k) \int_{\Sigma} \omega_0 \leq \int_{F_{r,k}(\Sigma)} \omega_k \leq 2 i_{g_k}^{-2}(p_k) \int_{\Sigma} \omega_0 \leq 2k^{-2} \int_{\Sigma} \omega_0 < 1,$$

for $k \gg 1$. Since $[F_{r,k}(\Sigma)] \in H_2(M_k, \mathbb{Z})$ and $[\omega_k] \in H^2(M_k, \mathbb{Z})$, we obtain

$$\int_{F_{r,k}(\Sigma)} \omega_k \in \mathbb{Z},$$

which is a contradiction. Hence $\omega_E|_{\bar{S}} \equiv 0$ and $\omega_0|_S \equiv 0$, which implies that S is a lagrangian submanifold (X, ω_0) by combining Lemma 3.1.

Since $\bar{S}$ is a lagrangian linear subspace of $(\mathbb{C}^n, \omega_E)$, there is a $\theta_0 \in \mathbb{R}$ such that $\text{Im} e^{\sqrt{-1}\theta_0} \Omega_E|_{\bar{S}} = 0$. This implies that $\bar{S}$ is a special lagrangian linear subspace of phase θ_0 in $(\mathbb{C}^n, \omega_E, \Omega_E)$. Thus S is a special lagrangian submanifold of phase θ_0 in (X, ω_0, Ω_0) , i.e.

$$\omega_0|_S = 0, \quad \text{Im} e^{\sqrt{-1}\theta_0} \Omega_0|_S = 0.$$

□

Lemma 3.3. For $k \gg 1$, $[F_{r,k}^* \tilde{\omega}_k|_S] = 0$ in $H^2(S, \mathbb{R})$.

Proof. By the smooth convergence of $\tilde{\omega}_k$ and Lemma 3.2,

$$\lim_{k \rightarrow \infty} \int_{F_{r,k}(A)} \tilde{\omega}_k = \int_A \omega_0 = 0,$$

for any cycle $A \in H_2(S, \mathbb{Z})$. For $k \gg 1$, we have

$$|\int_{F_{r,k}(A)} \omega_k| = i_{g_k}^2(p_k) |\int_{F_{r,k}(A)} \tilde{\omega}_k| < \frac{1}{2k^2} < 1.$$

Since $[F_{r,k}(A)] \in H_2(M_k, \mathbb{Z})$ and $[\omega_k] \in H^2(M_k, \mathbb{Z})$, we obtain $\int_{F_{r,k}(A)} \omega_k \in \mathbb{Z}$. This implies that $|\int_{F_{r,k}(A)} \omega_k| = 0$, and we obtain the conclusion

$$\int_A F_{r,k}^* \tilde{\omega}_k = 0.$$

□

Let $\tilde{X}$ be the total space of the pull-back $\pi_h^* \nu(S)$ of the normal bundle. Note that we can identify the zero section of $\pi_h^* \nu(S)$ with $\tilde{S}$, and the covering π_h extends to a finite covering $\pi : \tilde{X} \rightarrow X$ of X , i.e. $\tilde{S} = \pi^{-1}(S) \subset \tilde{X}$, and $\pi|_{\tilde{S}} = \pi_h$. The fundamental group $\pi_1(\tilde{S}) \cong \pi_1(\tilde{X})$ is isomorphic to the lattice Λ , $\pi_1(\tilde{X})$ is a normal subgroup of $\pi_1(X) = \pi_1(S)$, and the covering group $\Gamma \cong \pi_1(S)/\pi_1(\tilde{S}) = \pi_1(X)/\pi_1(\tilde{X})$. Note that $\pi_1(\tilde{X})$ (resp. $\pi_1(X)$) acts on $\mathbb{C}^n$ preserving g_E , ω_E and Ω_E , $\tilde{S}$ is invariant, $\tilde{X} = \mathbb{C}^n/\pi_1(\tilde{X})$ (resp. $X = \mathbb{C}^n/\pi_1(X)$), and $\tilde{S} = \tilde{S}/\pi_1(\tilde{S}) = \tilde{S}/\Lambda$ (resp. $S = \tilde{S}/\pi_1(S)$).

Proposition 3.4. *Let $\bar{S}^\perp$ be the orthogonal complement of $\bar{S}$ in $\mathbb{C}^n$, i.e. $\mathbb{C}^n = \bar{S} \oplus \bar{S}^\perp$, and $g_E(v, w) = 0$, for any $v \in \bar{S}$ and $w \in \bar{S}^\perp$. Then*

- i) $(\tilde{X}, \pi^* g_0)$ is isometric to $(T^n \times \bar{S}^\perp, h + h_E)$, where $T^n = \tilde{S}/\Lambda = \tilde{S}$, $h_E = g_E|_{\bar{S}^\perp}$, and h is the standard flat metric on T^n induced by $g_E|_{\bar{S}}$.
- ii) The action of Γ on $\tilde{X}$ is a product action, i.e. there are Γ -actions on T^n and $\bar{S}^\perp$ such that $\gamma \cdot (x, y) = (\gamma \cdot x, \gamma \cdot y)$ for any $\gamma \in \Gamma$, $x \in T^n$ and $y \in \bar{S}^\perp$. Furthermore, $T^n \times \{0\}$ is Γ -invariant, and $S = \pi(T^n \times \{0\}) = (T^n \times \{0\})/\Gamma$.
- iii)

$$\pi^* \omega_0|_{T^n \times \{y\}} \equiv 0, \quad \text{and} \quad \pi^* \text{Im} e^{\sqrt{-1}\theta_0} \Omega_0|_{T^n \times \{y\}} \equiv 0,$$

for any $y \in \bar{S}^\perp$, and a constant $\theta_0 \in \mathbb{R}$.

Proof. We choose coordinates $x_1, \dots, x_n$ on $\bar{S}$ and $y_1, \dots, y_n$ on $\bar{S}^\perp$ such that

$$g_E = \sum (dx_j^2 + dy_j^2), \quad \omega_E = \sum dx_j \wedge dy_j, \quad e^{\sqrt{-1}\theta_0} \Omega_E = \bigwedge_{j=1}^n (dx_j + \sqrt{-1} dy_j).$$

If $\mathcal{G}$ is a subgroup of the fundamental group $\pi_1(X) = \pi_1(S)$, then $\mathcal{G}$ acts on $\mathbb{C}^n$ preserving g_E , ω_E and Ω_E , and $\bar{S}$ is a invariant subspace. For any $\gamma \in \mathcal{G}$, we have $\gamma \cdot (v + w) = G_\gamma(v + w) + b_\gamma$, where $G_\gamma \in U(\mathbb{C}^n)$, $b_\gamma \in \bar{S}$, $v \in \bar{S}$ and $w \in \bar{S}^\perp$. Since $\bar{S}$ is invariant, we obtain then $G_\gamma(v + w) = A_\gamma v + B_\gamma w + C_\gamma w$ where $A_\gamma \in SO(\bar{S})$, $B_\gamma \in SO(\bar{S}^\perp)$, and $C_\gamma \in \text{Hom}(\bar{S}^\perp, \bar{S})$. Moreover, $G_\gamma \in SO(\mathbb{R}^{2n})$ implies $C_\gamma = 0$. Since $\omega_E(G_\gamma(v + w), G_\gamma(v + w)) = \omega_E(v + w, v + w)$, we have $B_\gamma = A_\gamma^{-1, T} = A_\gamma$, and $\gamma \cdot (v + w) = A_\gamma(v + w) + b_\gamma$. Thus $\pi_1(\tilde{X}) \cong \Lambda$ acts on $\mathbb{C}^n$ given by $\gamma \cdot (v + w) = v + w + b_\gamma$, $b_\gamma \in \Lambda$, for any $v \in \bar{S}$ and $w \in \bar{S}^\perp$. This implies that $\tilde{X} = \mathbb{C}^n/\pi_1(\tilde{X}) \cong \bar{S}/\Lambda \times \bar{S}^\perp = \tilde{S} \times \bar{S}^\perp$, and $\pi^* g_0 = h + h_E$ where $h_E = g_E|_{\bar{S}^\perp}$, and h is the standard flat metric on $\tilde{S}$ induced by $g_E|_{\bar{S}}$.

The $\pi_1(X)$ -action on $\mathbb{C}^n$ descends to a Γ -action on $\tilde{X}$, which is a product action since the $\pi_1(X)$ -action is so. Moreover, $\tilde{S} \times \{0\}$ is a invariant set as $\bar{S} \times \{0\}$ is invariant under the $\pi_1(X)$ -action. If we denote the quotient map $\mathbf{q}_1 : \mathbb{C}^n \rightarrow \mathbb{C}^n/\Lambda = \tilde{X}$, then

$\bar{\pi} = \pi \circ \mathbf{q}_1$, $g_E = \mathbf{q}_1^* \pi^* g_0$, $\omega_E = \mathbf{q}_1^* \pi^* \omega_0$, and $\Omega_E = \mathbf{q}_1^* \pi^* \Omega_0$. Since $\omega_E|_{\bar{S} \times \{y\}} = 0$ and $e^{\sqrt{-1}\theta_0} \Omega_E|_{\bar{S} \times \{y\}} = 0$ for $y \in \bar{S}^\perp$, we obtain that

$$\pi^* \omega_0|_{T^n \times \{y\}} \equiv 0, \quad \text{and} \quad \pi^* \text{Im} e^{\sqrt{-1}\theta_0} \Omega_0|_{T^n \times \{y\}} \equiv 0,$$

for a constant $\theta_0 \in \mathbb{R}$. $\square$

Remark 3.5. The coordinates $x_1, \dots, x_n$ on $\bar{S}$ in the proof of this proposition induce parallel 1-forms $dx_1, \dots, dx_n$ on $(\tilde{S}, h)$ which are pointwise linear independent, i.e. $dx_1, \dots, dx_n$ is a global parallel frame field. Under the coordinates $y_1, \dots, y_n$ on $\bar{S}^\perp$, we have these formulas

$$\pi^* g_0 = \sum (dx_j^2 + dy_j^2), \quad \pi^* \omega_0 = \sum dx_j \wedge dy_j, \quad e^{\sqrt{-1}\theta_0} \pi^* \Omega_0 = \bigwedge_{j=1}^n (dx_j + \sqrt{-1} dy_j).$$

Remark 3.6. The natural projection $f_0 : \tilde{X} \longrightarrow \bar{S}^\perp$ is equivariant under the Γ actions on $\tilde{X}$ and $\bar{S}^\perp$. For any $y \in \bar{S}^\perp$, $f_0^{-1}(y) = \tilde{S} \times \{y\}$, and f_0 is a special lagrangian fibration on $(\tilde{X}, \pi^* \omega_0, e^{\sqrt{-1}\theta_0} \pi^* \Omega_0)$, i.e. $\dim_{\mathbb{R}} f_0^{-1}(y) = n$,

$$\pi^* \omega_0|_{f_0^{-1}(y)} \equiv 0, \quad e^{\sqrt{-1}\theta_0} \pi^* \Omega_0|_{f_0^{-1}(y)} \equiv 0.$$

4. LOCAL SPECIAL LAGRANGIAN FIBRATIONS

In this section, we study the deformation of special lagrangian fibrations under the convergence of Calabi-Yau metrics. Let $(Y, \omega, g, J, \Omega)$ be a complete flat Calabi-Yau n -manifold.

Condition 4.1. *Assume that*

- i) $Y = T^n \times \mathbb{R}^n$, $g = h + h_E$, and the natural projection $f : Y \longrightarrow \mathbb{R}^n$ is a special lagrangian fibration of (Y, ω, Ω) , where $T^n = \mathbb{R}^n / \Lambda$ is a torus, Λ is a lattice in $\mathbb{R}^n$, h_E is the standard Euclidean metric on $\mathbb{R}^n$, and h is the standard flat metric induced by h_E .
- ii) We assume that there are parallel 1-forms $dx_1, \dots, dx_n$ on (T^n, h) , which are pointwise linear independent, and coordinates $y_1, \dots, y_n$ on $\mathbb{R}^n$ such that

$$g = h + h_E = \sum (dx_j^2 + dy_j^2), \quad \omega = \sum dx_j \wedge dy_j, \quad \Omega = \bigwedge_{j=1}^n (dx_j + \sqrt{-1} dy_j).$$

- iii) There is a family of Calabi-Yau structures $(\omega_k, g_k, J_k, \Omega_k)$ converging to (ω, g, J, Ω) in the C^∞ -sense on $Y_{2r} = T^n \times B_{h_E}(0, 2r)$ for a $r \gg 1$, where $B_{h_E}(0, 2r) = \{y \in \mathbb{R}^n \mid \|y\|_{h_E} < 2r\}$. Moreover, $\omega_k \in [\omega] \in H^2(Y_{2r}, \mathbb{R})$.
- vi) There is a finite group Γ acting on Y_{2r} preserving $\omega_k, g_k, \Omega_k, \omega, g, \Omega$, and $T^n \times \{0\}$ is a invariant set. The Γ -action is a product action on $T^n \times B_{h_E}(0, 2r)$. The natural projection $f : Y \longrightarrow \mathbb{R}^n$ is Γ -equivariant.

The goal of this section is to construct equivariant special lagrangian fibrations on $(Y_r, \omega_k, \Omega_k)$ for $k \gg 1$.

Denote $L = T^n \times \{0\}$, which is a special lagrangian submanifold of (Y, ω, Ω) , i.e. $\omega|_L = 0$ and $\text{Im} \Omega|_L = 0$. Note that we can identify Y with the total space of the normal bundle $\nu(L)$ by the exponential map from $\nu(L)$ to Y , $\exp_{L,g} : (x, \sum_j y_j \frac{\partial}{\partial y_j}) \mapsto (x, y)$ where $x \in L$ and $y = (y_1, \dots, y_n)$. There is a canonical bundle isomorphism from $\nu(L)$

to the cotangent bundle T^*L given by $v \mapsto \iota(v)\omega$ where $v \in \nu_x(L)$. Thus we can identify Y with the total space of T^*L by the map

$$(1) \quad (x, y) \mapsto (x, \iota(\sum_j y_j \frac{\partial}{\partial y_j})\omega) = (x, \sum_j y_j dx_j),$$

where $x \in L$ and $y = (y_1, \dots, y_n) \in \mathbb{R}^n$. We do not distinguish Y with T^*L in this section for convenience. For a 1-form σ on L , and a $y \in \mathbb{R}^n$, which can be regarded as a 1-form from above,

$$L(y, \sigma) = \{(x, y + \sigma(x)) | x \in L\}$$

denotes the graph of $y + \sigma$, i.e. $y = \sum y_j dx_j$, $\sigma = \sum \sigma_j dx_j$, and

$$L(y, \sigma) = \{(x, y_1 + \sigma_1(x), \dots, y_n + \sigma_n(x)) | x \in L\}.$$

There are two constants $a_k > 0$ and $\theta_k \in \mathbb{R}$, for any k , such that

$$\int_L \Omega_k = a_k e^{-\sqrt{-1}\theta_k} \int_L \Omega_0 = a_k e^{-\sqrt{-1}\theta_k} \int_L \text{Re}\Omega_0,$$

$\lim_{k \rightarrow \infty} a_k = 1$ and $\lim_{k \rightarrow \infty} \theta_k = 0$ by the smooth convergence of Ω_k . There are real 1-forms α_k and complex value $(n-1)$ -forms β_k such that

$$\omega_k = \omega_0 - d\alpha_k, \quad \Omega_k = a_k e^{-\sqrt{-1}\theta_k} (\Omega_0 + d\beta_k)$$

by $\omega_k \in [\omega] \in H^2(Y_{2r}, \mathbb{R})$. By the smooth convergence of ω_k and Ω_k ,

$$(2) \quad \lim_{k \rightarrow \infty} \|d\alpha_k\|_{C^2(Y_{2r}, g)} = \lim_{k \rightarrow \infty} \|d\beta_k\|_{C^2(Y_{2r}, g)} = 0.$$

Define a diffeomorphism $\Pi : L \rightarrow L(y, \sigma)$ by $x \mapsto (x, y + \sigma(x))$ for a $y \in \mathbb{R}^n$ and a 1-form σ on L . If

$$(3) \quad \mathfrak{F}_k(y, \sigma) = (-\Pi^* \omega_k|_{L(y, \sigma)}, *_h a_k^{-1} \Pi^* \text{Im} e^{\sqrt{-1}\theta_k} \Omega_k|_{L(y, \sigma)}),$$

where $*_h$ is the Hodge star operator on (L, h) , then $L(y, \sigma)$ is a special lagrangian submanifold of (Y, ω_k, Ω_k) of phase θ_k if and only if

$$\mathfrak{F}_k(y, \sigma) = 0.$$

A straightforward calculation (c.f. [34]) gives

$$(4) \quad \mathfrak{F}_k(y, \sigma) = (d\sigma + \Pi^* d\alpha_k|_{L(y, \sigma)}, *_h d *_h \sigma + *_h \Pi^* d\text{Im} \beta_k|_{L(y, \sigma)}).$$

We denote $\Omega^j(L)$ the space of j -forms on L , and define two Banach spaces $\mathfrak{B}_1 = C^{1, \alpha}(d\Omega^0(L) \oplus d^{*h} \Omega^2(L))$ and $\mathfrak{B}_2 = C^{0, \alpha}(d\Omega^1(L) \oplus d^{*h} \Omega^1(L))$. Then $\mathfrak{F}_k$ defines a smooth map $\mathfrak{F}_k : \mathcal{U}(r) \rightarrow \mathfrak{B}_2$ for any k , where $\mathcal{U}(r) = \{\|y\|_{h_E} + \|\sigma\|_{C^{1, \alpha}(L, h)} < 2r | (y, \sigma) \in \mathbb{R}^n \times \mathfrak{B}_1\}$.

Lemma 4.2. *For any $y \in B_{h_E}(0, 2r)$,*

$$\|\mathfrak{F}_k(y, 0)\|_{C^{0, \alpha}(L, h)} \leq C \|(d\alpha_k, d\beta_k)\|_{C^{1, \alpha}(Y_{2r}, g)},$$

for a constant C independent of k .

Proof. Since

$$\mathfrak{F}_k(y, 0) = (\Pi^* d\alpha_k|_{L(y, 0)}, *_h \Pi^* d\text{Im} \beta_k|_{L(y, 0)}),$$

we obtain the conclusion by straightforward calculations. $\square$

The differentials of $\mathfrak{F}_k(y, \sigma)$ are

$$(5) \quad D_\sigma \mathfrak{F}_k(y, \sigma) \dot{\sigma} = (d\dot{\sigma}, *_h d *_h \dot{\sigma}) + (D_\sigma(\Pi^* d\alpha_k|_{L(y, \sigma)}) \dot{\sigma}, *_h D_\sigma(\Pi^* d\text{Im}\beta_k|_{L(y, \sigma)}) \dot{\sigma}),$$

$$(6) \quad \begin{aligned} D_y \mathfrak{F}_k(y, \sigma) \dot{y} &= (D_y(\Pi^* d\alpha_k|_{L(y, \sigma)}) \dot{y}, *_h D_y(\Pi^* d\text{Im}\beta_k|_{L(y, \sigma)}) \dot{y}) \\ \text{and } D_\sigma \mathfrak{F}_k(y, \sigma)(\dot{y} + \dot{\sigma}) &= D_\sigma \mathfrak{F}_k(y, \sigma) \dot{\sigma} + D_y \mathfrak{F}_k(y, \sigma) \dot{y}. \end{aligned}$$

Under the frame field $dx_1, \dots, dx_n$ and coordinates $y_1, \dots, y_n$,

$$d\alpha_k = \sum_{ij} (\alpha_{k,ij} dx_i \wedge dx_j + \alpha_{k,i(n+j)} dx_i \wedge dy_j + \alpha_{k,(n+i)(n+j)} dy_i \wedge dy_j).$$

The differential is

$$\begin{aligned} D(\Pi^* d\alpha_k|_{L(y, \sigma)})(\dot{y} + \dot{\sigma}) &= \sum_{ijl} \left(\frac{\partial \alpha_{k,ij}}{\partial y_l} (\dot{y}_l + \dot{\sigma}_l) dx_i \wedge dx_j + \alpha_{k,i(n+j)} dx_i \wedge d\dot{\sigma}_j \right. \\ &\quad \left. + \frac{\partial \alpha_{k,i(n+j)}}{\partial y_l} (\dot{y}_l + \dot{\sigma}_l) dx_i \wedge d\sigma_j + \alpha_{k,(n+i)(n+j)} d\sigma_i \wedge d\dot{\sigma}_j \right. \\ &\quad \left. + \frac{\partial \alpha_{k,(n+i)(n+j)}}{\partial y_l} (\dot{y}_l + \dot{\sigma}_l) d\sigma_i \wedge d\sigma_j \right). \end{aligned}$$

We obtain

$$(7) \quad \begin{aligned} \|D_\sigma(d\alpha_k|_L)\dot{\sigma}\|_{C^{0,\alpha}(L,h)} &\leq C \|d\alpha_k\|_{C^{1,\alpha}(Y_{2r,g})} \|\dot{\sigma}\|_{C^{1,\alpha}(L,h)}, \\ \|D_\sigma(\Pi^* d\alpha_k|_{L(y, \sigma)})\dot{\sigma}\|_{C^{0,\alpha}(L,h)} &\leq C \|d\alpha_k\|_{C^{1,\alpha}(Y_{2r,g})} \left(\sum_{l=0,1,2} \|\sigma\|_{C^{1,\alpha}(L,h)}^l \right) \|\dot{\sigma}\|_{C^{1,\alpha}(L,h)}, \\ \|D_y(\Pi^* d\alpha_k|_{L(y, \sigma)})\dot{y}\|_{C^{0,\alpha}(L,h)} &\leq C \|d\alpha_k\|_{C^{1,\alpha}(Y_{2r,g})} \left(\sum_{l=0,1,2} \|\sigma\|_{C^{1,\alpha}(L,h)}^l \right) \|\dot{y}\|_{h_E}, \end{aligned}$$

for a constant C independent of k . The same argument gives

$$(8) \quad \begin{aligned} \|D_\sigma(d\text{Im}\beta_k|_L)\dot{\sigma}\|_{C^{0,\alpha}(L,h)} &\leq C \|d\beta_k\|_{C^{1,\alpha}(Y_{2r,g})} \|\dot{\sigma}\|_{C^{1,\alpha}(L,h)}, \\ \|D_\sigma(\Pi^* d\text{Im}\beta_k|_{L(y, \sigma)})\dot{\sigma}\|_{C^{0,\alpha}(L,h)} &\leq C \|d\beta_k\|_{C^{1,\alpha}(Y_{2r,g})} \left(\sum_{l=0,1,\dots,n} \|\sigma\|_{C^{1,\alpha}(L,h)}^l \right) \|\dot{\sigma}\|_{C^{1,\alpha}(L,h)}, \\ \|D_y(\Pi^* d\text{Im}\beta_k|_{L(y, \sigma)})\dot{y}\|_{C^{0,\alpha}(L,h)} &\leq C \|d\beta_k\|_{C^{1,\alpha}(Y_{2r,g})} \left(\sum_{l=0,1,\dots,n} \|\sigma\|_{C^{1,\alpha}(L,h)}^l \right) \|\dot{y}\|_{h_E}. \end{aligned}$$

Lemma 4.3. *The operator $D_\sigma \mathfrak{F}_k(0, 0)$ is invertible for $k \gg 1$, and*

$$\|D_\sigma \mathfrak{F}_k(0, 0)^{-1}\| \leq \overline{C},$$

for a constant $\overline{C} > 0$ independent of k .

Proof. Note that

$$D_\sigma \mathfrak{F}_k(0, 0) \dot{\sigma} = (d\dot{\sigma}, *_h d *_h \dot{\sigma}) + (D_\sigma(d\alpha_k|_L) \dot{\sigma}, *_h D_\sigma(d\text{Im}\beta_k|_L) \dot{\sigma}) = (\mathcal{D} + V_k) \dot{\sigma},$$

where $\mathcal{D} = d - *_h d *_h$ is the restriction of the Hodge Dirac operator $d + d^{*h}$ on the space of 1-forms, and, thus, is an elliptic operator of 1-order. By the standard elliptic estimate (c.f. Proposition 1.5.2 in [29] and [25]), we have

$$\|\xi\|_{C^{1,\alpha}(L,h)} \leq C_S \|\mathcal{D}\xi\|_{C^{0,\alpha}(L,h)},$$

for any $\xi \in \mathfrak{B}_1$, and a constant C_S independent of k . Hence $\mathcal{D}$ is injective. From the definition of $\mathfrak{B}_2$, $\mathcal{D}$ is also surjective, which implies that $\mathcal{D}$ is invertible from $\mathfrak{B}_1$ to $\mathfrak{B}_2$. Moreover,

$$\|\mathcal{D}^{-1}\| \leq C_S.$$

By (7) and (8),

$$\|V_k\| \leq C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})} < \frac{1}{2C_S},$$

for $k \gg 1$, and, thus,

$$\|\mathcal{D}^{-1}V_k\| < \frac{1}{2}.$$

By the standard operator's theory (c.f. [42]), $D_\sigma \mathfrak{F}_k(0,0) = \mathcal{D} + V_k$ is invertible, and the inverse operator is defined by

$$D_\sigma \mathfrak{F}_k(0,0)^{-1} = \left(\sum_{j=0}^{\infty} (-1)^j (\mathcal{D}^{-1}V_k)^j \right) \mathcal{D}^{-1}.$$

We obtain

$$\|D_\sigma \mathfrak{F}_k(0,0)^{-1}\| \leq \left(\sum_{j=0}^{\infty} 2^{-j} \right) \|\mathcal{D}^{-1}\| \leq \overline{C},$$

for a constant $\overline{C} > 0$ independent of k . $\square$

Lemma 4.4. *For any $\delta_0 \ll 1$, there is a constant $k_0 \gg 1$ such that, if $\|y\|_{h_E} \leq \frac{3r}{2}$ and $\|\sigma\|_{C^{1,\alpha}(L,h)} \leq \delta_0$, and $k > k_0$, then*

$$\|D_\sigma \mathfrak{F}_k(y, \sigma) - D_\sigma \mathfrak{F}_k(0,0)\| \leq \frac{1}{2\overline{C}}.$$

Furthermore, $D_\sigma \mathfrak{F}_k(y, \sigma)$ is also invertible, and

$$\|D_\sigma \mathfrak{F}_k(y, \sigma)^{-1}\| \leq 2\overline{C}.$$

Proof. By (5),

$$\begin{aligned} (D_\sigma \mathfrak{F}_k(y, \sigma) - D_\sigma \mathfrak{F}_k(0,0))\dot{\sigma} &= ((D_\sigma(\Pi^* d\alpha_k|_{L(y,\sigma)}) - D_\sigma(d\alpha_k|_L))\dot{\sigma}, \\ &\quad *_h(D_\sigma(\Pi^* d\text{Im}\beta_k|_{L(y,\sigma)}) - D_\sigma(d\text{Im}\beta_k|_L))\dot{\sigma}). \end{aligned}$$

We can take a $k_0 \gg 1$ such that, for $k > k_0$,

$$\begin{aligned} \|D_\sigma \mathfrak{F}_k(y, \sigma) - D_\sigma \mathfrak{F}_k(0,0)\| &\leq 2C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})} \left(\sum_{l=0,1,\dots,n} \|\sigma\|_{C^{1,\alpha}(L,h)}^l \right) \\ &\leq 2C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})} n\delta_0 \\ &\leq \frac{1}{4\overline{C}}, \end{aligned}$$

by (2), (7) and (8). We obtain the first formula in the conclusion.

Note that $D_\sigma \mathfrak{F}_k(y, \sigma) = D_\sigma \mathfrak{F}_k(0,0) + (D_\sigma \mathfrak{F}_k(y, \sigma) - D_\sigma \mathfrak{F}_k(0,0))$, $D_\sigma \mathfrak{F}_k(0,0)$ is invertible, and $\|D_\sigma \mathfrak{F}_k(0,0)^{-1}\| \leq \overline{C}$. By the same arguments as in the proof of Lemma 4.3, and

$$\|D_\sigma \mathfrak{F}_k(0,0)^{-1}(D_\sigma \mathfrak{F}_k(y, \sigma) - D_\sigma \mathfrak{F}_k(0,0))\| \leq \frac{1}{2},$$

$D_\sigma \mathfrak{F}_k(y, \sigma)$ is also invertible, and

$$\|D_\sigma \mathfrak{F}_k(y, \sigma)^{-1}\| \leq \left(\sum_{j=0}^{\infty} 2^{-j} \right) \|D_\sigma \mathfrak{F}_k(0,0)^{-1}\| \leq 2\overline{C}.$$

$\square$

Lemma 4.5. *For a fixed $\delta < \delta_0$, there is a $k_1 > k_0$ such that, for any $y \in B_{h_E}(0, \frac{3r}{2})$ and $k > k_1$, there is a unique $\sigma_k(y) \in \mathfrak{B}_1$, such that*

$$\mathfrak{F}_k(y, \sigma_k(y)) = 0, \quad \|\sigma_k(y)\|_{C^{1,\alpha}(L,h)} \leq \delta,$$

which implies that $L(y, \sigma_k(y))$ is a special lagrangian submanifold of $(Y_{2r}, \omega_k, \Omega_k)$. Furthermore,

$$\|D\sigma_k(y)\| \leq 2n\delta\overline{C}C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})},$$

for a constant C independent of k .

Proof. Fix a $\delta < \delta_0$, there is a $k_1 > k_0$ such that, for $k > k_1$, and any $y \in B_{h_E}(0, 2r)$,

$$\|\mathfrak{F}_k(y, 0)\|_{C^{0,\alpha}(L,h)} \leq \frac{\delta}{4\overline{C}},$$

by Lemma 4.2. By Theorem 2.7, Lemma 4.3 and 4.4, for any $y \in B_{h_E}(0, \frac{3r}{2})$ and $k > k_1$, there is a unique $\sigma_k(y) \in \mathfrak{B}_1$ such that

$$(9) \quad \mathfrak{F}_k(y, \sigma_k(y)) = 0, \quad \|\sigma_k(y)\|_{C^{1,\alpha}(L,h)} \leq \delta,$$

which implies that $L(y, \sigma_k(y))$ is a special lagrangian submanifold of $(Y_{2r}, \omega_k, \Omega_k)$.

By (6) (7) and (8),

$$\begin{aligned} \|D_y \mathfrak{F}_k(y, \sigma_k)\| &\leq C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})} \left(\sum_{l=0,1,\dots,n} \|\sigma_k\|_{C^{1,\alpha}(L,h)}^l \right) \\ &\leq C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})} n\delta, \end{aligned}$$

for a constant C independent of k . By Theorem 2.7,

$$D\sigma_k(y)\dot{y} = -D_\sigma \mathfrak{F}_k(y, \sigma_k)^{-1} D_y \mathfrak{F}_k(y, \sigma_k)\dot{y}.$$

We obtain the conclusion from Lemma 4.4. $\square$

Proposition 4.6. *For $k \gg 1$, there is an open set $Y_{2r} \supset W_k \supset Y_r$ such that $(W_k, \omega_k, \Omega_k)$ admits a equivariant special lagrangian fibration $f_k : W_k \longrightarrow B_k$ of phase θ_k over $B_k \subset \mathbb{R}^n$, i.e. there is a Γ -action on B_k , f_k is a Γ -equivariant map, and f_k is a special lagrangian fibration of phase θ_k , i.e.*

$$\omega_k|_{f_k^{-1}(b)} \equiv 0, \quad \text{Im}e^{\sqrt{-1}\theta_k}\Omega_k|_{f_k^{-1}(b)} \equiv 0,$$

for any $b \in B_k$.

Proof. By Lemma 4.5, there is a unique C^1 -map

$$\sigma_k : B_{h_E}(0, \frac{3r}{2}) \longrightarrow C^{1,\alpha}(d\Omega^0(T^n) \oplus d^{*h}\Omega^2(T^n)), \quad \text{by } y \mapsto \sigma_k(y),$$

which satisfies

$$\mathfrak{F}_k(y, \sigma_k(y)) = 0, \quad \|\sigma_k(y)\|_{C^{1,\alpha}(L,h)} \leq \delta \ll 1,$$

$$\text{and } \|D\sigma_k(y)\| \leq 2n\delta\overline{C}C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})}.$$

This implies

$$\left| \frac{\partial \sigma_{k,j}(y)}{\partial y_i} \right| \leq 2n\delta\overline{C}C\|(d\alpha_k, d\beta_k)\|_{C^{1,\alpha}(Y_{2r,g})} \ll 1,$$

for $k \gg k_1 > 1$.

Define a map $\Psi_k : Y_{\frac{3}{2}r} \longrightarrow Y_{2r}$ by

$$\Psi_k : (x, y) \mapsto (x, y_1 + \sigma_{k,1}(y), \dots, y_n + \sigma_{k,n}(y)) = (x, y + \sigma_k(y)).$$

Note that the frame field $dx_1, \dots, dx_n$ induces local coordinates $x_1, \dots, x_n$ around any point on L , and the differential can be expressed as

$$d\Psi_k : (\dot{x}, \dot{y}) \mapsto (\dot{x}_j + \sum \frac{\partial \sigma_{k,j}(y)}{\partial x_i} \dot{x}_i, \dot{y}_j + \sum \frac{\partial \sigma_{k,j}(y)}{\partial y_i} \dot{y}_i)$$

under such local coordinates. Thus $d\Psi_k$ is an isomorphism when $k \gg 1$, which implies that Ψ_k is an immersion. Furthermore, for $y_1 \neq y_2 \in \mathbb{R}^n$,

$$\begin{aligned} \Psi_k(x, y_2) - \Psi_k(x, y_1) &= (x, \dots, \int_0^1 (1 + \frac{\partial \sigma_{k,j}((1-t)y_2 + ty_1)}{\partial y_j} dt)(y_{2,j} - y_{1,j}), \dots) \\ &\neq 0. \end{aligned}$$

Hence Ψ_k is an embedding.

Note that the Γ -action on $Y_{2r} = T^n \times B_{h_E}(0, 2r)$ preserves $\omega_k, g_k, \Omega_k, \omega, g, \Omega$, and is a product action on $T^n \times B_{h_E}(0, 2r)$, i.e. there are Γ -actions on T^n and $B_{h_E}(0, 2r)$ such that $\gamma \cdot (x, y) = (\gamma \cdot x, \gamma \cdot y)$ for any $\gamma \in \Gamma$, $x \in T^n$, and $y \in B_{h_E}(0, 2r)$. Under the identification map (1),

$$\begin{aligned} (\gamma \cdot x, \gamma \cdot y) &= (\gamma \cdot x, \iota(\gamma_* \sum_j y_j \frac{\partial}{\partial y_j} \omega)) = (\gamma \cdot x, \gamma^* \omega(\sum_j y_j \frac{\partial}{\partial y_j}, \gamma_*^{-1} \cdot)) \\ &= (\gamma \cdot x, \gamma^{-1,*} \sum_j y_j dx_j). \end{aligned}$$

Thus

$$\begin{aligned} \gamma \cdot L(y, \sigma_k(y)) &= \{(\gamma \cdot x, \gamma \cdot (y_1 + \sigma_{k,1}(y)(x), \dots, y_n + \sigma_{k,n}(y)(x))) | x \in T^n\} \\ &= \{(\gamma \cdot x, \gamma^{-1,*} \sum_j (y_j + \sigma_{k,j}(y)(x)) dx_j | x \in T^n\} \\ &= L(\gamma \cdot y, \gamma^{-1,*} \sigma_k(y)), \end{aligned}$$

for any $\gamma \in \Gamma$ and $y \in B_{h_E}(0, 2r)$. Since the Γ -action preserves ω_k and Ω_k , $L(\gamma \cdot y, \gamma^{-1,*} \sigma_k(y))$ are special lagrangian submanifolds. By the uniqueness of $\sigma_k(y)$, $\gamma^{-1,*} \sigma_k(y) = \sigma_k(\gamma \cdot y) \in C^{1,\alpha}(d\Omega^0(T^n) \oplus d^{*h} \Omega^2(T^n))$. Hence

$$\begin{aligned} \Psi_k(\gamma \cdot x, \gamma \cdot y) &= (\gamma \cdot x, \gamma \cdot y + \sigma_k(\gamma \cdot y)) \\ &= (\gamma \cdot x, \gamma^{-1,*} \sum_j (y_j + \sigma_{k,j}(y)) dx_j) \\ &= (\gamma \cdot x, \gamma \cdot (y_1 + \sigma_{k,1}(y), \dots, y_n + \sigma_{k,n}(y))) \\ &= \gamma \cdot \Psi_k(x, y), \end{aligned}$$

i.e. Ψ_k is a Γ -equivariant map.

We denote $\mathcal{P} : Y_{2r} \rightarrow B_{h_E}(0, 2r)$ the natural projection, $B_k = B_{h_E}(0, \frac{3}{2}r)$ and $W_k = \Psi_k(Y_{\frac{3}{2}r})$. Since the Γ -action on $B_{h_E}(0, 2r)$ preserves the metric h_E and 0, $B_k = B_{h_E}(0, \frac{3}{2}r)$ is invariant. By $\delta \ll 1 \ll r$, $W_k \supset Y_r$. Then $f_k = \mathcal{P} \circ \Psi_k^{-1} : W_k \rightarrow B_k$ is a Γ -equivariant special lagrangian fibration of $(W_k, \omega_k, \Omega_k)$ of phase θ_k . We obtain the conclusion. $\square$

5. PROOF OF THEOREM 1.1

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Assume that the conclusion is not true. Then, for any fixed $\sigma > 1$, there is a family of closed Ricci-flat Calabi-Yau n -manifolds $\{(M_k, \omega_k, J_k, g_k, \Omega_k)\}$ with $[\omega_k] \in H^2(M_k, \mathbb{Z})$, and $p_k \in M_k$ such that

- i) the injectivity radius and the sectional curvature

$$i_{g_k}(p_k) < \frac{1}{k}, \quad \sup_{B_{g_k}(p_k, 1)} |K_{g_k}| \leq 1,$$

- ii) $[\Omega_k|_{B_{g_k}(p_k, \sigma i_{g_k}(p_k))}] \neq 0$ in $H^n(B_{g_k}(p_k, \sigma i_{g_k}(p_k)), \mathbb{C})$.
 iii) for any open subset $W'_k \supset B_{g_k}(p_k, \sigma i_{g_k}(p_k))$, $(W'_k, \omega_k, \Omega_k)$ wouldn't admit special lagrangian fibrations.

If we denote $\tilde{\omega}_k = i_{g_k}^{-2}(p_k)\omega_k$, $\tilde{g}_k = i_{g_k}^{-2}(p_k)g_k$, and $\tilde{\Omega}_k = i_{g_k}^{-n}(p_k)\Omega_k$, then

$$i_{\tilde{g}_k}(p_k) = 1, \quad \sup_{B_{\tilde{g}_k}(p_k, k)} |K_{\tilde{g}_k}| \leq \frac{1}{k^2},$$

and $[\tilde{\Omega}_k|_{B_{\tilde{g}_k}(p_k, \sigma)}] \neq 0$ in $H^n(B_{\tilde{g}_k}(p_k, \sigma), \mathbb{C})$. By the Cheeger-Gromov's convergence theorem (c.f. Theorem 2.4), a subsequence of $(M_k, \tilde{\omega}_k, \tilde{g}_k, J_k, \tilde{\Omega}_k, p_k)$ converges to a complete flat Calabi-Yau n -manifold $(X, \omega_0, g_0, J_0, \Omega_0, p_0)$ in the C^∞ -sense, i.e. for any $r > \sigma$, there are embeddings $F_{r,k} : B_{g_0}(p_0, r) \rightarrow M_k$ such that $F_{r,k}(p_0) = p_k$, and $F_{r,k}^* \tilde{g}_k$ (resp. $F_{r,k}^* \tilde{\omega}_k$ and $F_{r,k}^* \tilde{\Omega}_k$) converges to g_0 (resp. ω_0 and Ω_0) in the C^∞ -sense. Furthermore, $i_{g_0}(p_0) = 1$. The soul theorem (c.f. [11], [35]) implies that there is a compact flat totally geodesic submanifold $S \subset X$, the soul, such that (X, g_0) is isometric to the total space of the normal bundle $\nu(S)$ with a metric induced by $g_0|_S$ and a natural flat connection.

By Proposition (3.4), there is a finite normal covering $\pi : \tilde{X} \rightarrow X$ with covering group Γ such that

- i) $(\tilde{X}, \pi^* g_0)$ is isometric to $(T^n \times \mathbb{R}^n, h + h_E)$, where $T^n = \mathbb{R}^n / \Lambda$, Λ is a lattice in $\mathbb{R}^n$, h_E is the standard Euclidean metric on $\mathbb{R}^n$, and h is the standard flat metric on T^n induced by h_E .
 ii) The action of Γ on $\tilde{X}$ is a product action, i.e. there are Γ -actions on T^n and $\mathbb{R}^n$ such that $\gamma \cdot (x, y) = (\gamma \cdot x, \gamma \cdot y)$ for any $x \in T^n$ and $y \in \mathbb{R}^n$. Furthermore, $T^n \times \{0\}$ is Γ -invariant, and $S = (T^n \times \{0\})/\Gamma$.
 iii)

$$\pi^* \omega_0|_{T^n \times \{y\}} \equiv 0, \quad \text{and} \quad \pi^* \text{Im} e^{\sqrt{-1}\theta_0} \Omega_0|_{T^n \times \{y\}} \equiv 0,$$

for any $y \in \mathbb{R}^n$, and a constant $\theta_0 \in \mathbb{R}$.

Note that the Γ -action on $\mathbb{R}^n$ preserves h_E , and, $B_{h_E}(0, \rho)$, which implies that $\tilde{X}_\rho = T^n \times B_{h_E}(0, \rho)$ are invariant, for any $\rho > 0$. Lemma (3.3) shows $[F_{r,k}^* \tilde{\omega}_k|_S] = 0$ in $H^2(S, \mathbb{R})$, for $k \gg 1$, which implies $[\pi^* F_{r,k}^* \tilde{\omega}_k|_{\tilde{X}_\rho}] = 0$ in $H^2(\tilde{X}_\rho, \mathbb{R})$, for any $\rho > 0$. By Remark (3.5), there are parallel 1-forms $dx_1, \dots, dx_n$ on (T^n, h) , which are pointwise linear independent, and coordinates $y_1, \dots, y_n$ on $\mathbb{R}^n$ such that

$$\pi^* g_0 = \sum (dx_j^2 + dy_j^2), \quad \pi^* \omega_0 = \sum dx_j \wedge dy_j, \quad e^{\sqrt{-1}\theta_0} \pi^* \Omega_0 = \bigwedge_{j=1}^n (dx_j + \sqrt{-1} dy_j).$$

Hence Condition 4.1 is satisfied.

Let $r > \rho \gg \sigma$ such that $B_{g_0}(p_0, \sigma) \subset \pi(\tilde{X}_\rho) \subset \pi(\tilde{X}_{2\rho}) \subset B_{g_0}(p_0, r)$. By Proposition (4.6), for $k \gg 1$, there is an open set $W_k \supset \tilde{X}_\rho$ such that $(W_k, \pi^* F_{r,k}^* \omega_k, \pi^* F_{r,k}^* \Omega_k)$ admits

a equivariant special lagrangian fibration $f_k : W_k \longrightarrow B_k$ of phase θ_k , where $B_k \subset \mathbb{R}^n$, i.e. there is a Γ -action on B_k , f_k is a Γ -equivariant map, and

$$\pi^* F_{r,k}^* \omega_k|_{f_k^{-1}(b)} \equiv 0, \quad \pi^* F_{r,k}^* \text{Im} e^{\sqrt{-1}\theta_k} \Omega_k|_{f_k^{-1}(b)} \equiv 0,$$

for any $b \in B_k$. Hence f_k induces a special lagrangian fibration $\bar{f}_k : \pi(W_k) \longrightarrow B_k/\Gamma$, which implies that $(F_{r,k} \circ \pi(W_k), \omega_k, \Omega_k)$ admits a special lagrangian fibration, and $F_{r,k} \circ \pi(W_k) \supset B_{g_k}(x_k, \sigma i_{g_k}(x_k))$. It is a contradiction. We obtain the conclusion. $\square$

6. ESTIMATES FOR INJECTIVITY RADIUS

In this section, we prove Theorem 1.5 (Corollary 6.1) and Theorem 1.6 (Theorem 6.3). The following estimate for injectivity radius is a direct consequence of Theorem 2.6.

Corollary 6.1. *Let (M, g) be a closed Riemannian manifold, Θ be a calibration n -form, and $p \in M$. Assume that the sectional curvature K_g satisfies*

$$\sup_{B_g(p, 2\pi)} K_g \leq 1,$$

and there is a submanifold L calibrated by Θ such that $\dim_{\mathbb{R}} L = n$, $p \in L$, and

$$\int_L \Theta < \frac{\pi}{2n} \varpi_{n-1},$$

where ϖ_{n-1} is the volume of S^{n-1} with the standard metric of constant curvature 1. Then the injectivity radius $i_g(p)$ of (M, g) at p satisfies that

$$i_g(p)^n \leq \frac{n\pi^{n-1}}{2^{n-1}\varpi_{n-1}} \int_L \Theta.$$

Proof of Corollary 6.1 and Theorem 1.5. By Theorem 2.6, we have

$$\text{Vol}_{h_1}(B_{h_1}(r)) \leq \text{Vol}_g(B_g(p, r) \cap L) \leq \text{Vol}_g(L) = \int_L \Theta,$$

for any $r \leq \min\{i_g(p), \frac{\pi}{2}\}$, where h_1 denotes the standard metric on S^n with constant curvature 1, and $B_{h_1}(r)$ denotes a metric r -ball in S^n . Since $h_1 = dr^2 + \sin^2 r h_{S^{n-1}}$ where $h_{S^{n-1}}$ is the standard metric on S^{n-1} with constant curvature 1, we obtain $\sin r \geq \frac{2}{\pi}r$, and

$$\frac{2^{n-1}}{n\pi^{n-1}} r^n \varpi_{n-1} \leq \int_0^r \sin^{n-1} r dr \varpi_{n-1} = \text{Vol}_{h_1}(B_{h_1}(r)) \leq \int_L \Theta.$$

If $i_g(p) \geq \frac{\pi}{2}$, by letting $r = \frac{\pi}{2}$, we obtain

$$\frac{\pi}{2n} \varpi_{n-1} \leq \int_L \Theta < \frac{\pi}{2n} \varpi_{n-1},$$

which is a contradiction. Thus $i_g(p) < \frac{\pi}{2}$. By letting $r = i_g(p)$, we obtain

$$i_g(p)^n \leq \frac{n\pi^{n-1}}{2^{n-1}\varpi_{n-1}} \int_L \Theta.$$

We obtain Theorem 1.5 by applying the above arguments to special lagrangian submanifolds in Ricci-flat Calabi-Yau manifolds. $\square$

An obvious application of Corollary 6.1 is to estimate injectivity radiuses by volumes of holomorphic submanifolds, which has independent interests.

Corollary 6.2. *Let (M, ω, J, g) be a closed Kähler m -manifold, and $p \in M$. Assume that the sectional curvature K_g satisfies*

$$\sup_M K_g \leq 1,$$

and there is a smooth holomorphic n -submanifold N such that $p \in N$, and

$$\int_N \omega^n < \frac{(n-1)! \pi}{2} \varpi_{n-1}.$$

Then the injectivity radius $i_g(p)$ of (M, g) at p satisfies that

$$i_g(p)^n \leq \frac{\pi^{n-1}}{(n-1)! 2^{n-1} \varpi_{n-1}} \int_N \omega^n.$$

By combining this corollary and the result in [11], there are F -structures of positive rank on the regions of Kähler manifolds with bounded curvature and fibred by holomorphic submanifolds with small volumes.

Finally, we estimate the volume of metric balls when they are fibred by calibrated submanifolds with small volume in the absence of bounded curvature of the ambient manifolds.

Theorem 6.3. *For any $n, m \in \mathbb{N}$ and any $\varepsilon > 0$, there is a constant $\delta = \delta(n, m, \varepsilon) > 0$ satisfying that: Assume that (M, g, Θ) is a closed Ricci-flat Einstein m -manifold with a calibration n -form Θ , $p \in M$, and there is a homology class $A \in H_n(M, \mathbb{Z})$ such that for any $x \in B_g(p, 1)$, there is a n -submanifold L_x calibrated by Θ passing x and presenting A , i.e. $x \in L_x$ and $[L_x] = A$. If*

$$\int_A \Theta < \delta,$$

then

$$\text{Vol}_g(B_g(p, 1)) \leq \varepsilon.$$

Proof of Theorem 6.3 and Theorem 1.6. Assume that the conclusion is not true. Then there is a family of closed Ricci-flat Einstein m -manifolds $\{(M_k, g_k, \Theta_k)\}$ with calibration n -forms Θ_k , $p_k \in M_k$, and there are homology class $A_k \in H_n(M_k, \mathbb{Z})$ such that

- i) for any $x \in B_{g_k}(p_k, 1)$, there is a n -submanifold $L_{k,x}$ calibrated by Θ_k passing x and presenting A_k , i.e. $x \in L_{k,x}$ and $[L_{k,x}] = A_k$,
- ii)

$$\int_{A_k} \Theta_k < \frac{1}{k}.$$

However,

$$\text{Vol}_{g_k}(B_{g_k}(p_k, 1)) \geq C,$$

for a constant $C > 0$ independent of k .

By Gromov pre-compactness theorem (Theorem 2.2), by passing to a subsequence, $\{(M_k, g_k, p_k)\}$ converges to a complete path metric space (Y, d_Y, p) in the pointed Gromov-Hausdorff sense. By Theorem 2.3, the Hausdorff dimension of (Y, d_Y, p) is m , and there is a closed subset $S_Y \subset Y$ of Hausdorff dimension smaller than $m-1$, i.e. $\dim_{\mathcal{H}} S_Y < m-1$, such that $Y \setminus S_Y$ is a smooth manifold, and d_Y is induced by a Ricci-flat Einstein metric g_∞ on $Y \setminus S_Y$. Furthermore, for any compact subset $D \subset Y \setminus S_Y$, there are embeddings $F_{k,D} : D \rightarrow M_k$ such that $F_{k,D}^* g_k$ converges to g_∞ in the C^∞ -sense.

We take D big enough such that $D \cap B_{d_Y}(p, \frac{1}{8})$ is not empty. For a $y \in \text{int} D \cap B_{d_Y}(p, \frac{1}{4})$, there is a $r > 0$ such that $B_{g_\infty}(y, 2r) \subset \text{int} D \cap B_{d_Y}(p, \frac{1}{4})$, and $B_{g_k}(F_{k,D}(y), r) \subset F_{k,D}(D) \cap$

$B_{g_k}(p_k, \frac{1}{2})$ when $k \gg 1$. By the smooth convergence of $F_{k,D}^* g_k$, the sectional curvatures K_{g_k} satisfy

$$\sup_{F_{k,D}(D)} |K_{g_k}| \leq C_D,$$

for a constant C_D depending on D , but independent of k . Let L_k be a n -submanifold calibrated by Θ_k passing $F_{k,D}(y)$ and presenting A_k , i.e. $F_{k,D}(y) \in L_k$ and $[L_k] = A_k$. By Bishop-Gromov comparison theorem, for any $\rho \leq r$,

$$\begin{aligned} \frac{\text{Vol}_{g_k}(B_{g_k}(F_{k,D}(y), \rho))}{\rho^m} &\geq \text{Vol}_{g_k}(B_{g_k}(F_{k,D}(y), 1)) \geq \text{Vol}_{g_k}(B_{g_k}(p_k, \frac{1}{2})) \\ &\geq \frac{1}{2^m} \text{Vol}_{g_k}(B_{g_k}(p_k, 1)) \geq \frac{C}{2^m}. \end{aligned}$$

Then there is a uniform lower bound $\iota > 0$ for injectivity radiuses $i_{g_k}(F_{k,D}(y))$ at $F_{k,D}(y)$ (c.f. [35]), i.e. $i_{g_k}(F_{k,D}(y)) \geq \iota > 0$ for $k \gg 1$. By Theorem 2.6, we have

$$\text{Vol}_{h_1}(B_{h_1}(\rho)) \leq \text{Vol}_{g_k}(B_{g_k}(F_{k,D}(y), \rho) \cap L_k) \leq \int_{L_k} \Theta_k = \int_{A_k} \Theta_k \leq \frac{1}{k},$$

where $\rho = \min\{\iota, r, \frac{\pi}{\sqrt{C_D}}\}$, h_1 denotes the standard metric on S^n with constant curvature C_D , and $B_{h_1}(\rho)$ denotes a metric ρ -ball in S^n . We obtain a contradiction when $k \gg 1$, and, thus, we obtain the conclusion.

We obtain Theorem 1.6 by applying the above arguments to special lagrangian submanifolds in Ricci-flat Calabi-Yau manifolds. $\square$

Remark 6.4. Let $\{(M_k, g_k, \Theta_k)\}$ be a family of closed Ricci-flat Einstein m -manifolds with calibration n -forms Θ_k , and $\{p_k\}$ be a sequence of points such that there are open subsets $W_k \supset B_{g_k}(p_k, 1)$ in M_k admitting calibrated fibrations $f_k : W_k \rightarrow B_k$ corresponding to Θ_k , i.e. for any $b_k \in B_k$, $f_k^{-1}(b_k)$ is a n -submanifold calibrated by Θ_k . If

$$\lim_{k \rightarrow \infty} \int_{f_k^{-1}(b_k)} \Theta_k = 0,$$

where $b_k \in B_k$, Theorem 6.3 implies that

$$\lim_{k \rightarrow \infty} \text{Vol}_{g_k}(B_{g_k}(p_k, 1)) = 0,$$

and, by passing to a subsequence, $\{(M_k, g_k)\}$ converges to a path metric space of lower Hausdorff dimension in the pointed Gromov-Hausdorff sense.

Remark 6.5. Theorem 6.3 can be applied to G_2 -manifolds and $Spin(7)$ -manifolds since they are Ricci-flat Einstein manifolds (c.f. [29]).

REFERENCES

- [1] B.S.Acharya, *On mirror symmetry for manifolds of exceptional holonomy*, in *Mirror symmetry, vector bundles and lagrangian submanifolds*, Studies in Advanced Math. 23, (2001), 1-14.
- [2] M.T.Anderson, *Ricci curvature bounds and Einstein metrics on compact manifolds*, J. Amer. Math. Soc. 2 (1989), 455-490.
- [3] M.T.Anderson, *Convergence and rigidity of manifolds under Ricci curvature bounds*, Invent. Math. 102 (1990), 429-445.
- [4] M.T.Anderson, *The L^2 structure of moduli spaces of Einstein metrics on 4-manifolds*, G.A.F.A. (1991), 231-251.
- [5] J.Cheeger, *Degeneration of Einstein metrics and metrics with special holonomy*, in *Surveys in differential geometry VIII*, 29-73.

- [6] J.Cheeger, T.H.Colding, *On the structure of space with Ricci curvature bounded below I*, Jour. of Diff. Geom. 46(1997), 406-480.
- [7] J.Cheeger, T.H.Colding, *On the structure of space with Ricci curvature bounded below II*, Jour. of Diff. Geom. 52(1999), 13-35.
- [8] J.Cheeger, T.H.Colding, G.Tian, *On the singularities of spaces with bounded Ricci curvature*, Geom.Funct.Anal. Vol.12 (2002), 873-914.
- [9] J.Cheeger, K.Fukaya, M.Gromov, *Nilpotent structures and invariant metrics on collapsed manifolds*, Journal of the American Mathematical Society, 5 (1992), 327-372.
- [10] J.Cheeger, M.Gromov, *Collapsing Riemannian manifolds while keeping their curvature bound I*, J. Differ. Geom. 23 (1986), 309-364.
- [11] J.Cheeger, M.Gromov, *Collapsing Riemannian Manifolds while keeping their curvature bounded II*, J.Diff.Geom. 32 (1990), 269-298.
- [12] J.Cheeger, G.Tian, *Anti-self-duality of curvature and degeneration of metrics with special holonomy*, Commun. Math. Phys. 255 (2005), 391-417.
- [13] J.Cheeger, G.Tian, *Curvature and injectivity radius estimates for Einstein 4-manifolds*, Journal of the American Mathematical Society, 19 (2006), 487-525.
- [14] J.Duistermaat, *On global action-angle coordinates*, Comm. Pure Applied Math., 33 (1980), 687-706.
- [15] K.Fukaya, *Hausdorff convergence of Riemannian manifolds and its application*, Advance Studies in Pure Mathematics, 18 (1990), 143-234.
- [16] K.Fukaya, *Multivalued Morse theory, asymptotic analysis and mirror asymmetry*, Proceedings of Symposia in Pure Mathematics, 73 (2005), 205-278.
- [17] R.E.Green, H.Wu, *Lipschitz converges of Riemannian manifolds*, Pacific J. Math. 131 (1988), 119-141.
- [18] M.Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Birkhäuser 1999.
- [19] E.Goldstein, *A construction of new families of minimal lagrangian submanifolds via torus action*, J. Diff. Geom. 58 (2001), 233-261.
- [20] E.Goldstein, *Calibrated fibrations*, Comm. Anal. Geom. 10 (2002), 127-150.
- [21] M.Gross, *Examples of special lagrangian fibrations*, in Symplectic Geometry and Mirror Symmetry, World Scientific Singapore, (2001), 81-109.
- [22] M.Gross, *Special lagrangian fibrations II-Geometry*, Surveys in Differential Geometry: Differential geometry inspired by string theory, International Press, (1999), 341-404.
- [23] M.Gross, P.M.H.Wilson, *Mirror symmetry via 3-tori for a class of Calabi-Yau threefolds*, Math. Ann. 309 (1997), 505-531.
- [24] M.Gross, P.M.H.Wilson, *Large complex structure limits of K3 surfaces*, J. Diff. Geom. 55 (2000), 475-546.
- [25] D.Gilbarg, N.S.Trudinger, *Elliptic partial differential equations of second two*, Springer 1983.
- [26] S.Gukov, S.T.Yau, E.Zaslow, *Duality and fibrations on G_2 manifolds*, Turkish Journal of Mathematics, 27 (2003), 61-97.
- [27] R.Harvey, H.B.Lawson, *Calibrated geometries*, Acta Math., 148 (1982), 47-157.
- [28] N.Hitchin, *The moduli space of special lagrangian submanifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 25 (1997), 503-515.
- [29] D.D.Joyce, *Compact manifolds with special holonomy*, Oxford University Press, 2000.
- [30] D.D.Joyce, *Singularities of special lagrangian fibrations and the SYZ conjecture*, Comm.Anal.Geom. 11 (2003), 859-907.
- [31] D.D.Joyce, *Lectures on Calabi-Yau and special Lagrangian geometry*, math.DG/0108088.
- [32] M.Kontsevich, Y.Soielman, *Homological mirror symmetry and torus fibrations*, in *Symplectic geometry and mirror symmetry*, World Sci. Publishing, (2001), 203-263.
- [33] P.Lu, *Kähler-Einstein metrics on Kummer threefold and special lagrangian tori*, Comm. Anal. Geom. 7 (1999), 787-806.
- [34] R.C.Mclean, *Deformation of calibrated submanifolds*, Comm. Anal. Geom. 6 (1998), 705-747.
- [35] P.Petersen, *Riemannian Geometry*, Springer, 1997.
- [36] W.D.Ruan, *On the convergence and collapsing of Kähler metrics*, J. Differ. Geom. 52 (1999), 1-40.
- [37] W.D.Ruan, *Generalized special Lagrangian torus fibration for Calabi-Yau hypersurfaces in toric varieties. I*, Commun. Contemp. Math. 9 (2007), no. 2, 201-216.
- [38] W.D.Ruan, *Generalized special Lagrangian torus fibrations for Calabi-Yau hypersurfaces in toric varieties. II*, in *Mirror symmetry. V*, Amer. Math. Soc., Providence, RI, 2006. 457-477.
- [39] W.D.Ruan, *Generalized special Lagrangian fibration for Calabi-Yau hypersurfaces in toric varieties III: The smooth fibres*, arXiv:math/0309450.

- [40] W.D.Ruan, *H-minimal Lagrangian fibrations in Kahler manifolds and minimal Lagrangian vanishing tori in Kahler-Einstein manifolds*, arXiv:math/0309177 .
- [41] W.D.Ruan, Y.G.Zhang, *Convergence of Calabi-Yau manifolds*, arXiv:0905.3424.
- [42] S.Salur, *Deformations of special lagrangian submanifolds*, Commu. Cont. Math. Vol.2, 3 (2000), 365-372.
- [43] A.Strominger, S.T.Yau, E.Zaslow, *Mirror symmetry is T-duality*, Nucl. Phys. B479, (1996), 243-259.
- [44] V.Tosatti, *Limits of Calabi-Yau metrics when the Kähler class degenerates*, arXiv:0710.4571, to appear in J.Eur.Math.Soc. 2009.
- [45] R.P.Thomas, S.T. Yau, *Special Lagrangians, stable bundles and mean curvature flow*, Comm. Anal. Geom. 10 no. 5 (2002), 1075-1113.
- [46] S.T.Yau, *On the Ricci curvature of a compact Kähler manifold and complex Monge-Ampere equation I*, Comm. Pure Appl. Math. 31 (1978), 339-411.
- [47] S.T.Yau, *Einstein manifolds with zero Ricci curvature*, in *Lectures on Einstein manifolds*, International Press, (1999), 1-14.

DEPARTMENT OF MATHEMATICS, CAPITAL NORMAL UNIVERSITY, BEIJING, P.R.CHINA
E-mail address: yuguangzhang76@yahoo.com